
MEDICAL CASES OF DR.HUSSEIN, LEBANON

TheraNat SA Contact: courrier@theranat.com Tel 0041219223518 www.theranat.com



- 41 years old patient
- Work injury with cut of extensor tendons of left hand
- Patient wasn't able to close his fist



 MIL therapy was applied locally on the wound and on the anterior surface of wrist joint



- After 15 sessions of MIL therapy the hand regained its normal functions
- The patient was able to close his fist completely



- 94 years old woman diabetic, with petrochanteric fracture
- The X-ray shows the line of fracture



 After 15 sessions of MIL therapy calcification started to appear around the fracture



29-7-2003



- •64 years old woman
- Diabetic
- •Old burn



After 6 days




After 9 days



After 9 days



After 12 days



- •17 years old patient
- Tetra-pelagic caused by car accident

Bed Sour of the tetra-pelagic patient



Before treatment



After treatment

MIL therapy acting on a bed soure





MIL THERAPY IN...

MIL therapy in pediatrics





MIL Therapy in Treating Tennis Elbow



MIL in Post-Surgery Treatment



MILT in Treating Urology Complications




MIL Therapy in Dermatology

Female Cheek Diffuse Redness



3 Weeks After 3rd Treatment

Pre Treatment



Female Cheek Diffuse Redness



Pre Treatment



Number of treatments: 6 2 weeks apart

Female Cheek Facial Telangiectasia



Pre Treatment




Number of treatments: 3 weeks apart

Female Cheek facial Telangiectasia



Pre Treatment



Number of treatments: 3 2 weeks apart

Male Cheek facial Telangiectasia



Female Lip Venous Lake



Pre Treatment

Number of treatments: 1 1 month after treatment

Female Left Media Ankle telangiectasia



Pre Treatment



Number of treatments: 1 1 month after treatment

After Treatment

COPYRIGHT TheraNat SA

Female Back of Knee telangiectasia



MIL Therapy Results

in Other Specialities

Stomatitis

Herbal tinctures and medicinal solutions have been used to treat stomatitis but they take a lot of time.

As a rule, such treatments produce visible results on the seventh through tenth day. When MIL was applied, we sometimes observed the same effect right after the first treatment.

Full rehabilitation would occur on the third through fifth day.





COPYRIGHT TheraNat SA

Clinical Cases

- Московский НИИ педиатрии и детской хирургии МЗ России
- Московская Детская клиническая больница № 9 им.Г.Н.Сперанского



Quantum treatment of the skin autograft at the burn surface.

The skin autograft:





Without quantum treatment


Clinical Cases

- Московский НИИ педиатрии и детской хирургии МЗ России
- Московская Детская клиническая больница № 9 им.Г.Н.Сперанского

Cystic cystitis.


Grade 4 vesicoureteral reflux.

Preparation for ureter reimplantation.











Cystic cystitis resolved after 10 procedures.

Russian Oncological Scientific Center

Post-radiation skin reactions.

Positive effects have been achieved in all cases of treating post-radiation skin reactions.

A comparison of the time needed for the complete disappearance of local manifestations in children who underwent magneto-infrared-laser therapy with historical data has shown that LILR helped to shorten recovery time by 28%.





Clinical Cases

Research Center of Laser Medicine, Moscow

Quantum Therapy of Soft Tissue Purulent Wounds



Before treatment



After 7 sessions of quantum therapy



At 20 days following the start of therapy

Clinical Cases

Subacute cytomegalovirus infection provoked total alopecia.

Follicles were actually absent in the head skin before treatment (the «polished skin» symptom).



After 5 courses of combined quantum treatment. «Live» follicles and soft unevenly pigmented hair appeared.



After 11 treatment courses. The follow-up period was 11 months. New hair was healthy, glossy and evenly colored.